


Eating too much salt can lead to hypertension that will increase the risks of cardiovascular disease and stroke.







Hypertension

 Children aged 4 to 6 are recommended to take not more than 1200mg of sodium a day. (about half teaspoon of salt)*


* Data source: Dietary Reference Intakes for Chinese (2013 edition), Chinese Nutrition Society

Children aged 4 to 6 should take not more than 400mg of sodium per meal (based on 3 meals a day),

which is approximately equivalent to one of the following:









- Sodium is naturally present in some food ingredients such as meat, fish and eggs. Since the naturally present sodium counts towards the sodium intake per meal, the actual amount of seasonings used should be even less.
- When marinating or cooking with multiple seasonings, the amount used for each should be reduced.

Example

Based on preparation for 30 children, the amount of seasonings used per meal should not more than:

- 2 tablespoons of salt or
- about 210ml of soy sauce or
- about 340ml of oyster sauce or
- about 1L of ketchup

Note:

1 teaspoon \approx 5g or ml; 1 tablespoon \approx 15g or ml

Ways to reduce salt intake



Cook with natural and fresh spices

E.g. spring onion, garlic, ginger, coriander, herbs, a trace amount of pepper

Cook with fresh fruit

E.g. stir-fried meat with dragon fruit, steamed meat patty with apple cubes, stir-fried chicken fillets with fresh pineapple




Make sauce with fresh fruit and vegetables

E.g. fresh orange juice, fresh tomato sauce, spinach sauce, pumpkin sauce, sweet corn sauce



Avoid using high-sodium seasonings and ready-made sauces

E.g. fermented bean curd, MSG, chicken powder, teriyaki sauce



Use less salt

Minimise the use of salt in seasoning or cooking









